arXiv:cond-mat/0010252v1 [cond-mat.soft] 18 Oct 2000

Collective excitations of a two-dimensional interacting Bose gas in anti-trap and linear external potentials

D. V. Fil*
Institute for Single Crystals, National Academy of Sciences of Ukraine, Lenin av. 60, Kharkov 61001, Ukraine

S. I. Shevchenko
B. I. Verkin Institute for Low Temperature Physics and Engineering, National Academy of Sciences of Ukraine, Lenin av. 47, Kharkov 61164, Ukraine

We present a method of finding approximate analytical solutions for the spectra and eigenvectors of collective modes in a two-dimensional system of interacting bosons subjected to a linear external potential or the potential of a special form $u(x,y) = \mu - u\cosh^2 x/l$, where μ is the chemical potential. The eigenvalue problem is solved analytically for an artificial model allowing the unbounded density of the particles. The spectra of collective modes are calculated numerically for the stripe, the rare density valley and the edge geometry and compared with the analytical results. It is shown that the energies of the modes localized at the rare density region and at the edge are well approximated by the analytical expressions. We discuss Bose-Einstein condensation (BEC) in the systems under investigations at $T \neq 0$ and find that in case of a finite number of the particles the regime of BEC can be realized, whereas the condensate disappears in the thermodynamic limit.

PACS numbers: 03.75.Fi, 05.30.Jp

I. INTRODUCTION

The observation of BEC in alkali metal vapors [1–3] confined in a trap has opened the opportunities to investigate experimentally this macroscopic quantum phenomenon. That outstanding discovery has stimulated theoretical studies of BEC in inhomogeneous Bose gases subjected to external fields (see [4] for review). One of the problems under extensive investigation is the influence of an external potential on properties of low-dimensional Bose systems. It is known that an external potential may cause BEC in 2D ideal Bose gases at nonzero temperatures [5–7]. This problem was discussed in detail by Mullin [8]. Mullin has presented rigorous arguments why the Hohenberg theorem [9] ruling out BEC in 2D does not apply to the systems considered in [5–7].

Recently, the possibility of BEC in interacting 2D Bose gases has been investigated. Petrov, Holzmann and Shlyapnikov [10] considered a 2D Bose system confined in a harmonic oscillator potential. They have concluded that two regimes can be realized: a true condensate and a quasicondensate with fluctuating phase. In the mean field approximation the properties of such a system have been studied by Bayindir and Tanatar [11]. They demonstrate the similarity in thermodynamic behavior of ideal 2D Bose systems in a harmonic trap and those with weak interactions and a finite number of the particles. On the other hand, Mullin [8,12] argues the absence of BEC in trapped interacting Bose gases in 2D in the thermodynamic limit.

For understanding the macroscopic quantum properties of interacting Bose systems the key point is the behavior of elementary excitations. In two dimensions thermally excited fluctuations destruct the condensate. Therefore, the information about stability of BEC can be gained from the study of the spectrum and structure of low-energy excitations.

The theory for elementary excitations of interacting Bose systems has been derived by Bogoliubov [13]. After the experiments [1–3] this theory has received a new interest. Stringari [14] has obtained an analytical solution for the collective mode spectrum of a 3D Bose gas in an isotropic harmonic trap in the Tomas-Fermi regime. The same problem for the anisotropic trap has been resolved in Ref. [15]. Numerical solutions of the Bogoliubov equations for the trap geometry have been obtained in Ref. [16–20]. The exact result for the energies of the harmonic trap breathing modes in 2D has been found by Pitaevskii and Rosch [21]. The collective mode spectrum in a spatially periodic potential has been investigated in Ref. [22].

Further understanding of the properties of inhomogeneous 2D Bose systems requires the study of the low-energy excitations in the potentials different from the mentioned above. It is important to investigate the potentials, for

---

[*] e-mail fil@isc.kharkov.com

which an analytical solution of the eigenvalue problem can be found. In this paper we analyze two such cases. The first case corresponds to the potential $u(x,y) = \mu - u\cosh^2 x/l$, where μ is the chemical potential. Since the external field pushes the particles away from the region $x = 0$, and a rare density valley is formed at $x = 0$, we call such a geometry an "anti-trap" one. The second potential considered is a linear one: $u(x,y) = \mu - \alpha x$. The Bose cloud subjected to that potential has an edge at $x = 0$ with a linear dependence of the density on the distance to the edge.

An approximate analytical solution of the eigenvalue problem is obtained by the following method. We formally imply the density is unbounded at $|x| \to \infty$ ($x \to +\infty$ for the linear potential). Then, analytical expressions for the energies and the eigenvectors of collective excitations can be derived. To establish the applicability of the model description to real objects the eigenvalue problem is also solved numerically. It is shown that the spectra of collective modes localized at the rare density region (in the anti-trap potential) and at the edge (in the linear potential) are well approximated by the analytical solution. Using the analytical solution we calculate the one particle density matrix and analyze the off-diagonal long range order. We conclude that thermal fluctuations do not destroy completely the long range order in the systems with a finite number of the particles. It means that the BEC regime is realized. It is not the case in the thermodynamic limit, where the density of the condensate approaches to zero at $T \neq 0$.

In Sec. II a general description of elementary excitations in inhomogeneous 2D Bose systems is presented. The collective mode spectra in the anti-trap and the linear potential are obtained in Sec. III and Sec. IV, correspondingly. The one particle density matrix is calculated in Sec. V, and the conclusions are given in Sec. VI.

II. COLLECTIVE EXCITATIONS OF INHOMOGENEOUS 2D BOSE GAS

We consider an interacting 2D Bose gas in the external potential $u(\mathbf{r})$. The Hamiltonian of the system can be written in the following form

$$H = \int d^2r\{\frac{\hbar^2}{2m}\nabla\ \hat{\psi}^{+}(\mathbf{r})\nabla\hat{\psi}(\mathbf{r}) + [u(\mathbf{r}) - \mu]\hat{\psi}^{+}(\mathbf{r})\hat{\psi}(\mathbf{r}) + \frac{1}{2}\gamma\hat{\psi}^{+}(\mathbf{r})\hat{\psi}^{+}(\mathbf{r})\hat{\psi}(\mathbf{r})\hat{\psi}(\mathbf{r})\}, \tag{1}$$

where $\hat{\psi}$ is the boson field operator, m, the boson mass. In Eq. (1) we use the effective potential of the atom-atom interaction of the form $V(\mathbf{r}-\mathbf{r}') = \gamma\delta(\mathbf{r}-\mathbf{r}')$. The strength γ is considered as a parameter. To describe the collective excitations we use an approach developed by one of the authors in Refs. [23–25]. We introduce the phase $\hat{\varphi}(\mathbf{r})$ and density $\hat{\rho}(\mathbf{r})$ operators. They are connected with the Bose field operator by the relation

$$\hat{\psi} = e^{i\hat{\varphi}}\sqrt{\hat{\rho}}. \tag{2}$$

The commutation relation for the new operators is found to be

$$\hat{\rho}(\mathbf{r})\hat{\varphi}(\mathbf{r}') - \hat{\varphi}(\mathbf{r}')\hat{\rho}(\mathbf{r}) = i\delta(\mathbf{r}-\mathbf{r}'). \tag{3}$$

Using Eq. (2) we rewrite the Hamiltonian (1) as

$$H = \int d^2r\boldsymbol{\{}\frac{\hbar^2}{2m}\{\sqrt{\hat{\rho}}(\nabla\hat{\varphi})^2\sqrt{\hat{\rho}} + (\nabla\sqrt{\hat{\rho}})^2 + i[(\nabla\sqrt{\hat{\rho}})(\nabla\hat{\varphi})\sqrt{\hat{\rho}} - \sqrt{\hat{\rho}}(\nabla\hat{\varphi})(\nabla\sqrt{\hat{\rho}})]\} + [u(\mathbf{r}) - \mu]\hat{\rho} + \frac{\gamma}{2}\hat{\rho}^2\boldsymbol{\}}. \tag{4}$$

The density operator can be decomposed as

$$\hat{\rho}(\mathbf{r}) = \rho_0(\mathbf{r}) + \hat{\rho}_1(\mathbf{r}), \tag{5}$$

where $\rho_0(\mathbf{r})$ is determined by the condition that the expansion of Eq. (4) into powers of $\hat{\rho}_1$ and $\nabla\hat{\varphi}$ does not contain the linear terms. It yields the equation

$$u(\mathbf{r}) - \mu + \gamma\rho_0(\mathbf{r}) - \frac{\hbar^2}{2m}\frac{1}{\sqrt{\rho_0(\mathbf{r})}}\nabla^2\sqrt{\rho_0(\mathbf{r})} = 0. \tag{6}$$

Eq. (6) coincides with the Gross-Pitaevskii equation for the time independent part of the condensate wave function [4]. Therefore, the function $\rho_0(\mathbf{r})$ can be understood as the density of the particles at $T = 0$. At low temperatures the fluctuations of the density and the current are assumed to be small and one can neglect the anharmonic terms. The quadratic part of the Hamiltonian reads as

$$H_2 = \int d^2r\boldsymbol{\{}\frac{1}{4}[2\sqrt{\rho_0}\hat{\varphi}\hat{T}(2\sqrt{\rho_0}\hat{\varphi}) + \frac{\hat{\rho}_1}{\sqrt{\rho_0}}\hat{G}(\frac{\hat{\rho}_1}{\sqrt{\rho_0}})] + i\frac{\hbar^2}{4m}\{(\nabla\hat{\rho}_1)(\nabla\hat{\varphi}) - (\nabla\hat{\varphi})(\nabla\hat{\rho}_1) - \frac{\nabla\rho_0}{\rho_0}[\hat{\rho}_1(\nabla\hat{\varphi}) - (\nabla\hat{\varphi})\hat{\rho}_1]\}\boldsymbol{\}}, \tag{7}$$

where

$$\hat{T} = -\frac{\hbar^2}{2m}[\nabla^2 - \frac{1}{\sqrt{\rho_0}}(\nabla^2\sqrt{\rho_0})], \tag{8}$$

$$\hat{G} = \hat{T} + 2\gamma\rho_0. \tag{9}$$

Let us rewrite the operators $\hat{\varphi}$ and $\hat{\rho}_1$ in the second quantization representation in terms of creation ($\hat{b}_\nu^+$) and annihilation ($\hat{b}_\nu$) operators of the elementary excitations, which should also be bosons in the system of Bose particles. It is convenient to write $\hat{\varphi}$ and $\hat{\rho}_1$ in the following form [24]:

$$\hat{\rho}_1 = \sqrt{\rho_0}\sum_\nu [F_\nu \hat{b}_\nu + F_\nu^* \hat{b}_\nu^+], \tag{10}$$

$$\hat{\varphi} = \frac{1}{2i\sqrt{\rho_0}}\sum_\nu [\Theta_\nu \hat{b}_\nu - \Theta_\nu^* \hat{b}_\nu^+]. \tag{11}$$

The representation (10), (11) should transform the Hamiltonian (7) to the diagonal form

$$H_2 = const + \sum_\nu (E_\nu + \frac{1}{2})\hat{b}_\nu^+ \hat{b}_\nu, \tag{12}$$

where E_ν is the energy of the excitation with the quantum number ν. The Hamiltonian (7) is reduced to the form (12), if the functions $F_\nu(r)$, $\Theta_\nu(r)$ satisfy the equations

$$\hat{T}\Theta_\nu = E_\nu F_\nu, \tag{13}$$

$$\hat{G}F_\nu = E_\nu \Theta_\nu, \tag{14}$$

and are normalized by the condition

$$\int d^2 r \Theta_\nu(\mathbf{r}) F_\nu^*(\mathbf{r}) = 1. \tag{15}$$

One can check that the operators $\hat{\rho}$, $\hat{\varphi}$ in the form (10), (11) with F_ν, Θ_ν determined by Eqs. (13), (14), and (15) satisfy the commutation relation (3).

One can see that under the substitution $F_\nu = u_\nu + v_\nu$, $\Theta_\nu = u_\nu - v_\nu$ Eqs. (13), (14) coincide with the Bogoliubov-de Gennes equations [4]. We note also a similarity between our approach and the one developed by Ho and Ma [26].

III. COLLECTIVE MODE SPECTRUM IN THE ANTI-TRAP GEOMETRY

In this section we consider the special choice of the external potential:

$$u(x, y) = \mu - u\cosh^2\frac{x}{l}. \tag{16}$$

We assume the parameters of the potential satisfy the inequality $u \gg \hbar^2/2ml^2$. Then, one can neglect the fourth term in the left-hand side of Eq. (6), that corresponds to the Tomas-Fermi approximation. At $T = 0$ the density of the particles is equal to

$$\rho_0(x, y) = \frac{u}{\gamma}\cosh^2\frac{x}{l}. \tag{17}$$

To derive the spectrum of low-energy excitations ($E \ll u$) one can omit the kinetic energy terms $\hat{T}$ in the operator $\hat{G}$. Then, Eqs. (13), (14) yield the following differential equation for the function Θ

$$2u\cosh^2\frac{x}{l}\,\hat{T}\Theta_\nu(\mathbf{r})=E_\nu^2\Theta_\nu(\mathbf{r}). \tag{18}$$

The function F is given by the expression

$$F_\nu(\mathbf{r})=\frac{E_\nu}{2u}\cosh^{-2}\frac{x}{l}\;\Theta_\nu(\mathbf{r}). \tag{19}$$

Taking into account that the system is uniform in the y direction the function Θ_ν can be presented in the form

$$\Theta_\nu(x,y)=\theta_\nu(x)e^{iky}, \tag{20}$$

where $\theta_\nu(x)$ satisfies the equation

$$-\frac{\hbar^2 u}{m}\cosh^2\frac{x}{l}\,[\frac{d^2\theta_\nu}{dx^2}-(k^2+l^{-2})\theta_\nu]=E_\nu^2\theta_\nu. \tag{21}$$

Eq. (21) coincides with the Shroedinger equation for the one particle problem in the potential $v(x)=\cosh^{-2}(x/l)$. Its solution given in Ref. [27] has the form

$$\theta_\nu(x)=\cosh^{-\eta}\frac{x}{l}\;w(\frac{1-\tanh\frac{x}{l}}{2}), \tag{22}$$

where $\eta=\sqrt{1+(kl)^2}$, and the function $w(z)$ is governed by a hypergeometric differential equation

$$z(1-z)w''+(\eta+1)(1-2z)w'-(\eta-s)(\eta+s+1)w=0. \tag{23}$$

In Eq. (23) the parameter s is determined by the equation $E_\nu^2=\omega_0^2 s(s+1)$, where $\omega_0=\sqrt{\hbar^2u/ml^2}$. To solve Eq. (23) one should specify the boundary conditions. We consider the system infinite in the x direction. Since the fluctuations of the current should remain finite, the quantity $d(\theta(x)/\sqrt{\rho_0(x)})/dx$ cannot be singular at $|x|\to\infty$. The solutions of Eq. (23) satisfying such a condition are the polynomials. Eq. (23) has the polynomial solutions at $s=\eta+n$, where $n=0,1,2,\ldots$. Therefore, the energy of the collective modes are equal to

$$E_{nk}=\omega_0\sqrt{[n+\sqrt{1+(kl)^2}][n+1+\sqrt{1+(kl)^2}]}. \tag{24}$$

The exact expression for the eigenfunction $\theta_{nk}(x)$ reads as

$$\theta_{nk}(x)=C_{nk}\cosh^{-\eta}\frac{x}{l}\;P_n^{\eta\eta}(\tanh\frac{x}{l}), \tag{25}$$

where $P_n^{\eta\eta}$ is the Jacobi polynomial. The normalization factor C_{nk} is calculated to be

$$C_{nk}=\left(\frac{u_0}{lL_yE_{nk}}\frac{n!(2n+2\eta+1)\Gamma(n+2\eta+1)}{2^{2\eta+1}\Gamma^2(n+\eta+1)}\right)^{1/2}. \tag{26}$$

In Eq. (26) Γ is the Gamma function, L_y, the size of the system in the y direction.

The answers obtained have two uncommon features. First, the spectrum does not contain an acoustic branch. Second, all the modes are localized near $x=0$. Such a peculiar behavior is connected with that the model contains an unphysical assumption. We fix the parameter l, while set an infinite size of the system in x direction. Thus, we allow the density of particles be unbounded at $|x|\to\infty$. The principal question has to be clarified is whether the results of the model (further we call it the artificial model) can be applied to real physical objects.

Let us consider two examples. The first one is a 2D Bose cloud of a stripe shape. We assume that there are two rigid walls at $x=\pm L_x$, and the potential well obeys the internal structure given at $|x|<L_x$ by Eq. (16). Since the particles cannot flow though the rigid walls, the gradient of the phase normal to the wall should be zero at $|x|=\pm L_x$. Therefore, the boundary conditions for the function $\theta(x)$ are

$$\frac{d}{dx}\left(\frac{\theta(x)}{\sqrt{\rho_0(x)}}\right)\Big|_{x=\pm L_x}=0. \tag{27}$$

The general solution of Eq. (21) can be written in the form

$$\theta(x) = \cosh\frac{x}{l}\ [A_1\varphi_1(x) + A_2\varphi_2(x)], \tag{28}$$

where

$$\varphi_1(x) = \cosh^{-\eta-1}\frac{x}{l}\ \ {}_2F_1(\eta - s, \eta + s + 1, \eta + 1, \frac{1 - \tanh\frac{x}{l}}{2}), \tag{29}$$

$$\varphi_2(x) = \cosh^{-\eta-1}\frac{x}{l}\ (1 - \tanh\frac{x}{l})^{-\eta}\ \ {}_2F_1(-s, s + 1, 1 - \eta, \frac{1 - \tanh\frac{x}{l}}{2}). \tag{30}$$

In Eqs. (29), (30) ${}_2F_1(x, y, z, u)$ is the hypergeometric function. We specify $1 - \eta \neq -1, -2, \ldots$. (If $1 - \eta$ is negative integer, another form of the general solution should be used.) Using Eqs. (27), (28) we get the dispersion equation

$$\varphi_1'(L_x)\varphi_2'(-L_x) - \varphi_1'(-L_x)\varphi_2'(L_x) = 0. \tag{31}$$

Its solutions can be obtained numerically. It is instructive to analyze the case of $L_x \gg l$ (at $L_x \approx l$ the density variation across the stripe is small). The solution of Eq. (31) at $L_x = 3l$ is shown in Fig. 1(a). For comparison, the spectrum given by the artificial model (Eq. (24)) is also presented in this figure. One can see that in the stripe system the lowest collective mode is an acoustic one at small k. Its velocity is equal to $\bar{c} = \sqrt{\gamma\bar{\rho}/m}$, where $\bar{\rho}$ is the average density of the particles. There is a discrepancy in behavior of the two families of the curves in Fig. 1(a) only at

$$E > \hbar\bar{c}k. \tag{32}$$

If the inequality opposite to (32) is satisfied, one can neglect the influence of the boundaries. In this case the collective mode spectrum is well approximated by Eq. (24).

To consider another example we specify the case of a Bose cloud infinite in the x direction with the density of the particles given by Eq. (17) at $|x| \leq L_x$ and the uniform density at $|x| \geq L_x$. There are localized and extended collective modes in such a system. The energies of the extended modes satisfy the inequality (32). The localized mode solution has the form $\theta(x) = A_3\exp(\kappa x)$ at $x \leq -L_x$, $\theta(x) = A_4\exp(-\kappa x)$ at $x \geq L_x$, where $\kappa = \sqrt{k^2 - E^2/\hbar^2\bar{c}^2}$. At $|x| \leq L_x$ it is given by Eq. (28). The boundary conditions are reduced to the continuity of $\theta(x)$ and $d(\theta(x)/\sqrt{\rho_0(x)})/dx$ at $x = \pm L_x$. The dispersion equation has the form

$$[\varphi_1'(-L_x) - \kappa\varphi_1(-L_x)][\varphi_2'(L_x) + \kappa\varphi_2(L_x)] - [\varphi_1'(L_x) + \kappa\varphi_1(L_x)][\varphi_2'(-L_x) - \kappa\varphi_2(-L_x)] = 0. \tag{33}$$

The solutions of Eq. (33) at $L_x = 3l$ are shown in Fig. 1(b). This figure illustrates that the analytical expression (24) yields a satisfactory approximation for the spectra of localized modes. From the numerical analysis we find that the coincidence is better for larger ratio L_x/l.

Thus, our artificial model, while is not free from some artifacts, seems quite useful for obtaining the analytical expressions for the spectra and eigenfunctions of collective modes for real physical objects.

IV. COLLECTIVE MODE SPECTRUM IN THE LINEAR POTENTIAL

In this section we consider the case of a linear external potential within the approach presented in Sec. III. This problem was studied preliminary in Ref. [24].

The external potential has the form

$$u(x, y) = \mu - \alpha x. \tag{34}$$

Let us consider the low-energy excitations in the Tomas-Fermi regime. In this regime the density of the particles is equal to $\rho_0(x, y) = \alpha x/\gamma$.

The equation for $\theta(x)$ is reduced to

$$-\frac{\hbar^2\alpha}{m}x[\frac{d^2\theta_\nu}{dx^2} + (\frac{1}{4x^2} - k^2)\theta_\nu] = E_\nu^2\theta_\nu. \tag{35}$$

Substituting

$$\theta_\nu(x) = \sqrt{2|k|x}e^{-|k|x}w(2|k|x) \tag{36}$$

into Eq. (35), we obtain the following equation for the function $w(z)$:

$$zw'' + (1-z)w' - \beta w = 0 \tag{37}$$

with

$$\beta = \frac{1}{2}\left(1 - \frac{E_\nu^2 m}{\hbar^2 |k| \alpha}\right). \tag{38}$$

At the beginning, we consider the artificial model. We assume the system is semi-infinite and the density is unbounded at $x \to \infty$. We are interested in the solutions finite at $x = 0$ and $x \to \infty$. Such solutions exist if $\beta = -n$ $(n = 0, 1, 2, \ldots)$. Using Eq. (38) we obtain the following analytical expression for the energies:

$$E_{nk} = \sqrt{\frac{\hbar^2 \alpha}{m}|k|(2n+1)}. \tag{39}$$

The function $\theta_\nu(x)$ reads as

$$\theta_{nk}(x) = C_{nk}\sqrt{2|k|x}e^{-|k|x} L_n(2|k|x), \tag{40}$$

where L_n is the Laguerre polynomial. The normalization factor is equal to

$$C_{nk} = \sqrt{\frac{\alpha}{L_y E_{nk}}}. \tag{41}$$

As in the case of the anti-trap potential the spectrum of the excitations does not contain an acoustic branch. It is connected with the unbounded density at $x \to \infty$ implied in the model.

Let us switch to the real systems and consider the stripe with a linear increase of the density across the stripe and a rigid wall at $x = L_x$. The solution of Eq. (35) finite at $x = 0$ has the form

$$\theta(x) = A\sqrt{x}e^{-|k|x} \ {}_1F_1(\beta, 1, 2|k|x), \tag{42}$$

where ${}_1F_1(x, y, z)$ is the Kummer confluent hypergeometric function. Using the boundary condition Eq. (27) at $x = L_x$, we obtain the following dispersion equation

$${}_1F_1(\beta, 1, 2|k|L_x) - 2\beta \ {}_1F_1(1+\beta, 2, 2|k|L_x) = 0. \tag{43}$$

Its solutions are shown in Fig. 2(a). The spectrum (39) is also presented in Fig. 2. In this figure the energy units $\epsilon_0 = \sqrt{\hbar^2\alpha/mL_x}$ are used. One can see that at $k > E/\hbar\bar{c}$ the spectrum is well approximated by the analytical expression (39), while at smaller k the difference is essential. In particular, the lowest mode transforms to the acoustic one.

To complete the picture we consider a semi-infinity system with a linear increase of the density at $0 \le x \le L_x$ and the uniform density at $x \ge L_x$. We are interested in the edge collective excitations of such a system. The dispersion equation for the edge modes is found to be

$$(\kappa - |k|) \ {}_1F_1(\beta, 1, 2|k|L_x) + 2|k|\beta \ {}_1F_1(1+\beta, 2, 2|k|L_x) = 0. \tag{44}$$

The solution of Eq. (44) is shown in Fig. 2(b). One can see that the spectrum of the edge excitation is very close to the dependence given by the analytical expression (39).

To conclude this section we discuss the conditions, when the low-energy description is valid. The Tomas-Fermi approximation is correct at

$$\gamma\rho_0(x) \gg \frac{\hbar^2}{2mx^2}, \tag{45}$$

when the fourth term in the left-hand side of Eq. (6) can be neglected. In two dimensions the parameter γ is connected with the radius of the interparticle interaction a by the relation [28]

$$\gamma = -\frac{4\pi\hbar^2}{m}\frac{1}{\ln(\rho_0 a^2)}. \tag{46}$$

Since Eq. (46) contains only the factor $\ln(\rho_0 a^2)^{-1}$ as a small parameter, the condition (45) is fulfilled at $x > x_0$, where x_0 is of order of the local average distance between the particles.

To omit the kinetic energy terms in the operator $\hat{G}$ we should consider the excitations with the energies

$$E < \gamma\rho_0(x_0) = \alpha x_0. \tag{47}$$

Since we neglect the solution of Eq. (37) singular at $kx \to 0$, the wave vector k should satisfy the inequality

$$kx_0 \ll 1. \tag{48}$$

Using Eqs. (39), (47), (48) we find the conditions on the wave vectors and the energies

$$k \ll \left(\frac{\alpha m}{\hbar^2}\right)^{1/3}, \tag{49}$$

$$E \ll E_m = \left(\frac{\hbar^2\alpha^2}{m}\right)^{1/3}. \tag{50}$$

Eqs. (49), (50) set the validity of the low-energy approximation. At finite L_x the energies have the scale of ϵ_0. Therefore, the description requires $\epsilon_0 \ll E_m$, that yields the condition on the size of the system in the x direction

$$L_x \gg \left(\frac{\hbar^2}{\alpha m}\right)^{1/3}. \tag{51}$$

V. OFF-DIAGONAL LONG RANGE ORDER

It is of interest to consider the possibility of BEC in 2D systems subjected to the anti-trap or linear potential. Let us analyze the behavior of the one-particle density matrix

$$W(\mathbf{r},\mathbf{r}') = \langle \hat{\Psi}^+(\mathbf{r})\hat{\Psi}(\mathbf{r}')\rangle. \tag{52}$$

If the function $W(\mathbf{r},\mathbf{r}')$ remains finite at $|\mathbf{r}-\mathbf{r}'| \to \infty$, the off-diagonal long range order is realized.

Thermal fluctuations destruct the macroscopic quantum coherence. The principal mechanism of the destruction is connected with the phase fluctuations. They are described by the operator $\hat{\varphi}$. Neglecting a small contribution of the density fluctuations we obtain the following expression for the function $W(\mathbf{r},\mathbf{r}')$:

$$W(\mathbf{r},\mathbf{r}') \approx \sqrt{\rho_0(\mathbf{r})\rho_0(\mathbf{r}')}\exp(\frac{-\Phi(\mathbf{r},\mathbf{r}')}{2}), \tag{53}$$

where

$$\Phi(\mathbf{r},\mathbf{r}') = \langle[\hat{\varphi}(\mathbf{r}) - \hat{\varphi}(\mathbf{r}')]^2\rangle. \tag{54}$$

To be more concrete, we analyze the dependence of $\Phi(r,r')$ on y, y' at $x = x'$. Taking into account Eq. (10) we get

$$\Phi(x,y;x,y') = \frac{1}{\rho_0(x)}\sum_{n=0}^{\infty}\frac{L_y}{2\pi}\int_{-\infty}^{+\infty} dk|\theta_{nk}(x)|^2[n_B(E_{nk}) + \frac{1}{2}][1-\cos k(y-y')], \tag{55}$$

where $n_B(E) = [\exp(E/T) - 1]^{-1}$.

Let us consider the case of the anti-trap potential. Since the maximum of the phase fluctuations is located at $x = 0$ (see. Eq. (25)), the phase coherence destruction begins at $x = 0$. Therefore, if the long range order survives at $x = 0$, it takes place in the whole system.

Using the analytical expressions (24), (25) for E_{nk} and $\theta_{nk}(x)$ we obtain the following asymptotic result:

$$\Phi(0,y,0,y') = \frac{T}{2\pi T_0(0)}\ln\frac{l}{\xi_T(0)} + \Phi_{\mathrm{zp}}, \tag{56}$$

where $T_0(x) = \hbar^2\rho_0(x)/2m$, $\xi_T(x) = \sqrt{\hbar^2\gamma\rho_0(x)/mT^2}$. In Eq. (56) the inequalities $l \gg \xi_T(0)$, $|y-y'| \gg l$ are implied. The term $\Phi_{\rm zp}$ describes a contribution of the zero-point fluctuations. This term determines the density of the condensate $n_0(x)$ at $T=0$. The temperature dependent part in the right-hand side of Eq. (56) does not depend on $|y-y'|$. It means that the condensate survives at nonzero temperatures. According to Eq. (56) the condensate density at $T \neq 0$ is connected with $n_0(0)$ by the expression

$$n_T(0) = n_0(0)\Big(\frac{\xi_T(0)}{l}\Big)^{\frac{T}{4\pi T_0(0)}}. \tag{57}$$

The result (57) is obtained within the artificial model. Nevertheless, one can apply it to the real objects, if in Eq. (55) the contribution of the modes with the energies (32) is negligible. For example, such a situation is realized for a Bose cloud of the shape of a square $L \times L$ with $L/l \gg 1$. Then, the answer (57) yields a satisfactory approximation for the density of the condensate. For the square geometry the total number of the particles N is connected with the parameter of the model by the relation

$$\frac{N}{L^2} = \frac{u}{2\gamma}(1 + \frac{l}{L}\sinh\frac{L}{l}). \tag{58}$$

It is natural to define the thermodynamic limit as $N \to \infty$, $L \to \infty$ at $N/L^2 = const.$ From Eq. (58) we also obtain $L/l = const.$ Therefore, in the thermodynamic limit the parameter l depends on the total number of the particles ($l \sim N^{1/2}$). From Eq. (57) we find that the density of the condensate decreases by the law $n_T(0) \sim N^{-b}$ ($b > 0$) under increase of N. It indicates the absence of true BEC in the thermodynamic limit. At finite N, as it follows from Eq. (57), the BEC regime can be realized.

Similar calculations for the linear potential yield

$$n_T(x) = n_0(x)\Big(\frac{\xi_T(x)}{x}\Big)^{\frac{T}{4\pi T_0(x)}}. \tag{59}$$

In Eq. (59) the inequalities $x \gg \xi_T(x)$, $|y-y'| \gg x$ are implied. For a Bose cloud of the shape of a rectangle $L_x \times L_y$ the thermodynamic limit corresponds to $N \to \infty$, $L_x \to \infty$, and $L_y \to \infty$ at $N/(L_xL_y) = const$ and $L_x/L_y = const.$ To analyze the density of the condensate at the same local density of the particles we should also fix the ratio x/L_x in Eq. (59). Then again $n_T(x)$ approaches to zero at $N \to \infty$ and there is no BEC in the thermodynamic limit.

VI. CONCLUSIONS

We have considered the low-energy collective excitations of the interacting inhomogeneous Bose gas in two-dimensions. We specify the two types of the external potential: the anti-trap one, in which the rare density valley is formed, and the linear potential, that models the simplest case of the edge geometry. We show that for the configurations considered an approximate analytical solution for the energies and the eigenvectors of the collective modes can be obtained. To derive the analytical solution we consider the artificial model, in which an unbounded density of the particles is allowed. Physically, such an approximation means that we suppress the fluctuations in the high-density regions. Our method yields the modes localized at the rare density valley or at the edge. Their spectra does not contain an acoustic branch. The modes lost in our approach can be obtained from the numerical solution of the eigenvalue problem. In this case we should specify the high-density profile. We present the results of the numerical analysis and compare them with the analytical solutions. We find that the spectra obtained numerically are well approximated by the analytical expressions in case of $E < \hbar k\sqrt{\gamma\bar{\rho}/m}$, where E is the energy of the excitation, k, the wave vector along the valley or the edge, $\bar{\rho}$, the total average density. It is important to note that in this case the spectrum of the collective modes is determined completely by the parameters of the external potential and does not depend on the interaction between bosons. Collective modes of the trapped Bose gas in the Tomas-Fermi regime demonstrate the same behavior [14].

The possibility of BEC in the systems under consideration has been investigated. The high density regions affect significantly on the character of the thermal fluctuations in the low density areas. If the total number of the particles is finite, the BEC regime is realized at nonzero temperatures. This conclusion is in agreement with the results of Ref. [10], where the BEC regime of a 2D Bose gas in the trapping potential has been described. The proper formulation of the thermodynamic limit for the systems considered shows that in this limit the true BEC does not occur, as it should follow from the general theorems [9,29,30].

ACKNOWLEDGMENTS

The authors would like to thank G. V. Shlyapnikov for helpful discussions. This work was supported by the INTAS Grant No. 97-0972.

[1] M. H. Anderson, J. R. Ensher, M. R. Matthew, C. E. Wieman, and E. A. Cornell, Science **269**, 198 (1995).
[2] C. C. Bradley, C. A. Sackett, J. J. Tolett, and R. G. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).
[3] K. B. Davis, M. O. Mewes, M. R. Andrews, N. J. van Druten, D. S. Durfee, D. M. Kurn, and W. Ketterle, Phys. Rev. Lett. **75**, 3969 (1995).
[4] F. Dalfovo, G. Giorgini, L. P. Pitaevskii, S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999).
[5] R. Masut, and W. J. Mullin, Am. J. Phys. **47**, 493 (1979).
[6] S. I. Shevchenko, Fiz. Nizk. Temp. **16**, 119 (1990) [Sov. J. Low. Temp. Phys. **16**, 64 (1990)].
[7] V. Bagnato and D. Kleppner, Phys. Rev. A **44**, 7439 (1991).
[8] W. J. Mullin, J. Low Temp. Phys. **106**, 615 (1997).
[9] P. C. Hohenberg, Phys. Rev. **158**, 383 (1967).
[10] D. S. Petrov, M. Holzmann, and G. V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000).
[11] M. Bayindir and B. Tanatar, Phys. Rev. A **58**, 3134 (1998).
[12] W. J. Mullin, J. Low Temp. Phys. **110**, 167 (1998).
[13] N. N. Bogoliubov, J. Phys. (USSR) **11**, 23 (1947).
[14] S. Stringari, Phys. Rev. Lett. **77**, 2360 (1996).
[15] P. Öhberg, E. L. Surkov, I. Tittonen, S. Stenholm, M. Wilkens, and G. V. Shlyapnikov, Phys. Rev. A **56**, R3346 (1997).
[16] P. A. Ruprecht, M. Edwards, K. Burnett, and C. W. Clark, Phys. Rev. A **54**, 4178 (1996); M. Edwards, P. A. Ruprecht, K. Burnett, R. J. Dodd, and C. W. Clark, Phys. Rev. Lett. **77**, 1671 (1996).
[17] D. A. W. Hutchinson, E. Zaremba, Phys. Rev A **57**, 1280 (1997); D. A. W. Hutchinson, E. Zaremba, and A. Griffin, Phys. Rev. Lett. **78**, 1842 (1997).
[18] K. G. Singh and D. S. Rokhsar, Phys. Rev. Lett. **77**, 1667 (1996).
[19] B. D. Eary, Phys. Rev. A **55**, 1147 (1997).
[20] L. You, W. Hoston, and M. Lewenstein, Phys. Rev. A **55**, 1581 (1997); V. M. Perez-Garcia, H. Michinel, J. I. Cirac, M. Lewenstein, and P. Zoller, Phys. Rev. Lett. **77**, 5320 (1996).
[21] L. P. Pitaevskii and A. Rosch, Phys. Rev. A **55** R853 (1997).
[22] Kirstine Berg-Sorensen and Klaus Molmer, Phys. Rev. A **58**, 1480 (1998).
[23] S. I. Shevchenko, Zh. Exp. Teor. Fiz. **100**, 1824 (1991) [Sov. Phys. JETP **73**, 1009 (1991)].
[24] S. I. Shevchenko, Fiz. Nizk. Temp. **18**, 328 (1992) [Sov. J. Low Temp. Phys. **18**, 223 (1992)].
[25] S. V. Terentjev and S. I. Shevchenko, Fiz. Nizk. Temp. **25**, 664 (1999) [Low Temp. Phys. **25**, 493 (1999)].
[26] Tin-Lun Ho and Michael Ma, J. Low Temp. Phys. **115**, 61, (1999).
[27] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics* (Butterworth-Heinemann, Oxford, 1999).
[28] V. N. Popov, *Functional Integrals in Quantum Field Theory and Statistical Physics* (D.Reidel Publishing, Dordrecht, 1983).
[29] O. Penrose and L. Onsager, Phys. Rev. **104**, 576 (1956).
[30] C. V. Chester, M. E. Fisher, and N. D. Mermin Phys. Rev. **185**, 760 (1969).



FIG. 1. Collective mode spectrum in the anti-trap potential for the stripe (a) and the rare density valley geometry (b). Solid curves show the numerical results; dashed curves, the result of Eq. (24).



FIG. 2. Collective mode spectrum in the linear potential for the stripe (a) and the semi-infinite system (b). Solid curves show the numerical results; dashed curves, the result of Eq. (39).